

National Kidney Foundation Of Michigan, Inc.



1169 Oak Valley Drive • Ann Arbor, Michigan 48108 • In State 800-482-1455 • 734-222-9800 • Fax 734-222-9801 • www.nkfm.org

House Health Policy Committee November 1, 2005 Supporting House Bill 5063

Chronic kidney disease (CKD) is a common progressive health problem. One in nine adults has CKD and most do not know it. It is the ninth leading cause of death in Michigan and is considered a growing silent epidemic in the United States. In 2004 there were 11,040 people on dialysis and 5,000 people living with a kidney transplant in Michigan. CKD can be prevented in many populations that are at high risk, and for those who already have CKD, progression towards kidney failure can be delayed. However, in most cases, CKD is not being detected early enough to initiate treatment regimens and reduce death and disability. Early identification (using GFR testing) is recommended.

Glomerular filtration rate (GFR) indicates the ability of the kidney's to filter and is an excellent measure of kidney function. It is calculated from the creatinine value in blood test results and is expressed as a percentage (%) of kidney function. A healthy GFR is 90 or above. A GFR of 15 or less usually indicates kidney failure, and the need for dialysis or kidney transplant to sustain life. If kidneys have already been damaged (GFR between 30-60) medications called ACE inhibitors and Angiotensin Receptor Blockers (ARB's) can protect the kidneys and keep them working for as long as possible.

HB 5063 would allow the Michigan Department of Community Health to perform "laboratory services, diagnostic services, early intervention services, and treatment for CKD" for its eligible Medicaid patients. This makes good health sense and good economic sense. Treatments for CKD costs much less than treatments for kidney failure! HB 5063 along with voluntary efforts aimed at physicians, health plans and medical laboratories can help reduce the number of people whose kidneys fail in Michigan.

The National Kidney Foundation of Michigan recommends that the House Health Policy Committee vote to support HB 5063.

Sally Joy **Public Policy Consultant** National Kidney Foundation of Michigan (kidney failure1985)



Chronic Kidney Disease & GFR: A Clinical Action Plan

American Journal of Kidney Disease, February 2000


Stage	Prevalence, U.S. adults	Description	GFR	Action
	20 million	At increased risk for chronic kidney disease, mostly people with diabetes, hypertension	≥90	Kidney disease screening
1	5.9 million	Kidney damage with normal GFR	≥90	Diagnosis & treatment to slow/stop progression
2	5.3 million	Kidney damage with declining GFR	60-89	Continue treatments to slow/stop progression
3	7.6 million	Moderate decline in GFR	30-59	Evaluate & treat complications
4	400,000	Severe decline in GFR	15-29	Prepare for dialysis or kidney transplant
5	300,000	Kidney failure	<15	Dialysis or kidney transplant necessary to maintain life
1-5	20 million	With chronic kidney disease		

CKD = chronic kidney disease

CVD = cardiovascular disease

GFR = glomerular filtration rate (% of kidney function - normal is 90-100).

patients in the Medicare & EGHP Pts, 2002-3 Figure p.34 Distribution of costs for CKD, CHF, & diabetic




CKD Size ? スの音でのの

> 四日 ひば

> > 120N

CKD costs ?



Population estimated from the 5 percent Medicare sample, & includes patients surviving all of 2002 with Medicare as primary payor. Diagnoses determined from claims in 2002. Patients with ESRD in the 5 percent sample are excluded. Costs are for calendar year 2003, with patients censored at development of ESRD.

USRDS 2005 ADR



October 31, 2005

Honorable Representative Gaffney, Representative Gleason and members of the House Health Policy Committee:

I am writing on behalf of the Gift of Life Minority Organ and Tissue Transplant Education Program (MOTTEP) to ask for your support of House Bill 5063.

According to the United Network for Organ Sharing, there are more than 83,000 individuals on the national transplant waiting list. Sixteen people die each day waiting for a life saving organ transplant. Minorities make up more than half of the national kidney transplant waiting list and, in Michigan, 46% of those currently waiting for a kidney transplant are African-American. Furthermore chronic kidney disease (CKD) is a prevalent health problem unknowingly affecting many people. It is the ninth leading cause of death in Michigan and in many cases could have been prevented if detected early enough to initiate treatment. House Bill 5063 will help reduce the number of people waiting for kidney transplants and save many lives by treating people before they enter end stage renal kidney failure.

Gift of Life MOTTEP is active in the metropolitan Detroit area. Our mission is to decrease the number of ethnic and minority Americans needing organ and tissue transplants, encourage healthy lifestyles and behavioral patterns and most importantly promote prevention through our "Love Yourself, Take Care of Yourself" message.

Saving lives and preventing disease is our goal and the goal of House Bill 5063. Your support is greatly appreciated.

Remonia Chapman
Director Gift of Life MOTTEP

2203 Platt Road, Ann Arbor, MI 48104

2515 579 427



LABORATORY REPORT

1215 E. Michigan Ave. Lansing, MI 48912 517.364.7800 • FAX 517.364.7810 800.884.2522

ATIENT

5176553775

CCOUNTSHARP, MARY P.

LANSING NEPHROLOGY GRP. 1717 E. MICH AVE, #A

LANSING


MI

48912

HISTORY NUMBER	REQU	JISITION NUMBER	3 O	ORDER STATUS	
181992	RM F	20506400	32 FI	NAL.	
COLLECTION DATE	C	OLLECTION TIME	R	EPORT DATE	
SEE BELOW	SE	E BELOW	0370	03/06/05	
AGE / D.O.B.	SEX	AREA	ROUTE	PAGE	
69Y 09/25/3	EF	01	04	1	

TEST DESCRIPTION	ON .	RESULT		EXPECTED RANGE	GE UNITS
ESTS ORDERED BY: OPY TO:	SHARP, MARY P. PATIENT COPY	MD			
Coll: 03/05/05 0					
COMPLETE BLOOD CO	est performed at	RM/Lab 800	E.Columbi	a Mason, MI	
WBC	20141	5.5		/ A 4 A A	14 4 600 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
RBC		3.52		4.0-12.0 3.50-5.55	K/CU MM
HGB		11.5	L	12.0-15.0	M/CU MM
HCT		32.5	L.	36.0-45.0	G/DL
MCV		92	I	80-100	% ————————————————————————————————————
MCH		3e. 7		27.0-33.0	CU MICR
MCHC		35. 4		31.0-37.0	MMCG %
FLATELET COUNT		257		150-400	K/CU MM
NEUTROPHILS		47.9	L	49.0-81.0	K/CO MM %
LYMPHOCYTES		30.8	1	14.0-41.0	/= "/u
MONOCYTES		9.5		0.0-11.0	/u "/ _n
EOSINOPHILS		10.9	Н	0.0-6.0	/n "/"
BASOPHILS		ō., 9	• •	0.0-3.0	"/"
		- " -		"" a "." C., a "."	, yu
RENAL FUNCTION PA	NEL				
CALCIUM		9.6		8.0-10.5	MG/DL
PHOSPHOROUS		4.7	Н	2.5-4.5	MG/DL
ALBUMIN		4.E		3.6-5.0	G/DL
BUN		29	Н	6-23	MG/DL
CREATININE		1.4		0.6-1.4	MG/DL
GLUCOSE		92		65-99	MG/DL
	NORMAL RANGE EFFE		1-04		
SODIUM		141		135-145	MEQ/L
POTASSIUM(K+)		4.8		3.5-4.9	MEQ/L
COE		25.0		20.0-30.0	MMOL/L
CHLORIDE		105		96-110	MEQ/L
ANION GAP		1 1		2-16	
CALCULATED GFR		39.12			ML/MIN
IF THE PATI	ENT IS AFRICAN-AM	MERICAN YOL	MUST MULT	TIPLY THE GFR E	3Y 1.18
	HRONIC KIDNEY DIS				
STAGE				ML/MIN/1.735Q.	M)
1	KIDNEY DAMAGE WOR ELEVATED GFR		> 5	90	
Ξ '	KIDNEY DAMAGE W DECREASED GFR	IITH MILD	60	-89	
3	MODERATE DECREA	SED GFR	30-	-59	
4	SEVERE DECREASE		15-		
	KIDNEY FAILURE	en out I i		OR DIALYSIS	
	A C 4 she bears track I V from	^	المالية ا	miv mrumiato	

EPATIC PANEL





Reg#: 18342763 Name: FERRITER, MAURICE J DOB: 06/07/1952 Sex: M Age: 52 Years User Name: DTRAMEL

More medadenacia			Pathology Handbook			
T = 1 - 7 - 4 O - 40	Order Test Name		Last Updated			
Order Test Code		Ordered:05/27/2005 12:05				
TACRO	CREATINE PHOSPHOKINASE					
СК			Updated:05/27/2005 13:28			
			30-240 IU/L			
Collected:05/27/2005 08:25 CREATINE PHOSPHOKINASE (CK)						
			Updated:05/27/2005 13:20			
	139		136-146 MEQ/L			
		Н	3.5-5.0 MEQ/L			
			99-111 MEQ/L			
CHLORIDE (CHLOR)			24-34 MEQ/L			
		11	8-20 MG/DL			
REA NITROGEN (UN)	1		0.9-1.3 MG/DL			
CREATININE (CREAT) GLUCOSE (GLUC)		П	73-110 MG/DL			
			8.6-10.2 MG/DL			
	CK HOSPHOKINASE (CK) BASIC SODIUM (SOD) POTASSIUM (POT) CHLORIDE (CHLOR CO2 (CO2) REA NITROGEN (UN) CREATININE (CREAT) GLUCOSE (GLUC CALCIUM (CAL	Order Test Code Order Test Name TACRO TACROLIMUS (FK506) CK CREATINE PHOSPHOKINASE HOSPHOKINASE (CK) 60 BASIQ Basic Metabolic Panel SODIUM (SOD) 139 POTASSIUM (POT) 5.1 CHLORIDE (CHLOR) 103 CO2 (CO2) 26 REA NITROGEN (UN) 73 CREATININE (CREAT) 3.0 GLUCOSE (GLUC) 110 CALCIUM (CAL) 9.2	Order Test Code Order Test Name TACRO TACROLIMUS (FK506) CK CREATINE PHOSPHOKINASE HOSPHOKINASE (CK) 60 BASIQ Basic Metabolic Panel SODIUM (SOD) 139 POTASSIUM (POT) 5.1 H CHLORIDE (CHLOR) 103 CO2 (CO2) 26 REA NITROGEN (UN) 73 H CREATININE (CREAT) 3.0 H CALCIUM (CAL) 9.2 TALCIUM (CAL)			

The MDRD formula for estimation of GFR (eGFR) was developed in a population of adults (>19 years old) with slowly-declining or stable reduced kidney function. The MDRD should not be used to predict eGFR in unstable patients or in children. A GFR estimate between 15 and 59 ml/min for >=3 months is classified as chronic

between 15 and 59 mi/min 100					
kidney disease (Stage 3 or 4).	28	L	> 59 ML/MIN		
AFRICAN AMERICAN EGFR (B-EGFR	/	Ī	> 59 MUMIN		
NON-AFRICAN AMERICAN EGFR (NB-EGFR	2		Updated:05/27/2005 13:03		
ACCN: 05-147-2011 CRCP	CBCP		Updated:05/27/2000-15:05		
Collected:05/27/2005 08:25					
MOV NORMAL RANGE: 8.7 to 11.6 11.	8.4		4.0-10.0 K/MM3		
WHITE BLOOD CELL COUNT (WBC	/	L	13.0-17.3 G/DL		
HEMOGLOBIN (HGE	//	L	39.0-50.2 %		
HEMATOCRIT (HCT	404		150-450 K/MM3		
PLATELET COUNT (PLT	/	L	4.50-5.90 M/MM3		
RED BLOOD CELL COUNT (RBC	7]	H	80.0-100.0 fl		
MEAN CORPUSCULAR VOLUME (MCV	/	```	25.0-35.0 pg		
MEAN CORPUSCULAR HGB (MCH	/		30.0-37.0 %		
MEAN CORPUSCULAR HGB CONCENTRN (MCHC	/	Н	11,5-15.5 %		
RED CELL DISTRIBUTION WIDTH (RDV			fl		
MEAN PLATELET VOLUME (MP)) 14:1)	L			
ACCN: 05-147-2011 AUTO DIFF	AUTO DIFF (5 PT FROM COL	JLTER)	Updated:05/27/2005 13:03		
Collected: 05/27/2005 08:25		***************************************	36.0-75.0 %		
NEUT % FROM COULTER (NEUT %	7/		20.0-50.0 %		
LYMPH % FROM COULTER (LYMPH %	7/1		3.0-10.0 %		
MONO % FROM COULTER (MONO 9	//		.0-4.0 %		
EOS % FROM COULTER (EOS %	3/1		.0-2.0 %		
BASO % FROM COULTER (BASO %	2/	<u> </u>			
ACCN: 05-147-2011 ABSOLT CNT	ABSOLUTE COUNTS		Updated:05/27/2005 13:03		
Collected:05/27/2005 08:25	4.7		1.4-7.5 K/MM3		
ABSOLUTE NEUTROPHIL COUNT (NEUT			0.8-5.0 K/MM3		
ABSOLUTE LYMPHOCYTE COUNT (LYMPH			0.1-1.0 K/MM3		
ABSOLUTE MONOCYTE COUNT (MONO			0.0-0.4 K/MM3		
ABSOLUTE EOSINOPHIL COUNT (EOS	7)		0.0-0.2 K/MM3		
ABSOLUTE BASOPHIL COUNT (BASO	τη . <u></u>	<u></u>			
ACCN: 05-147-2011 CHD	CORONARY HEART DISEAS	E PROFILE	Updated:05/2/12005 12:42		
Collected:05/27/2005 08:25		1			



For Immediate Release

Contacts:

Karen Glowacki/Ellie Schlam National Kidney Foundation 212-889-2210 x 171/143 kareng@kidney.org ellies@kidney.org

NEW DIAGNOSIS CODES FOR CHRONIC KIDNEY DISEASE TO BE BASED ON NATIONAL KIDNEY FOUNDATION'S K/DOQI GUIDELINES

NEW YORK, May 31, 2005 -Expanded codes and improved terminology will lead to better care of patients with chronic kidney disease (CKD). That is the opinion of the National Kidney Foundation (NKF), which developed the staging system for CKD soon to be included in the International Classification of Diseases, Ninth Revision Clinical Modification (ICD-9-CM). Beginning October 1, 2005, the new codes will allow medical professionals to clearly note the stage of kidney disease they diagnose and treat.

The staging system was developed by NKF through its Kidney Disease Outcomes Quality Initiative (KDOQI). Its 2002 clinical practice guideline on CKD classification defined CKD according to objective criteria irrespective of cause and classified it into five distinct stages based on the level of kidney function. These stages were used by the National Center for Health Statistics (NCHS) in this revision of the ICD-9-CM codes. It will now be easier for physicians to bill, keep data and plan treatment based on stage of disease.

"Until now, the majority of CKD patients were lumped into the chronic renal failure category, which was not very well defined. CKD has five stages, based on estimates of glomerular filtration rate, and the treatment plan for each of these stages becomes more detailed and specific as the disease progresses," said David Warnock, M.D., president of the NKF. "We are optimistic that by using the NKF-K/DOQI guidelines in conjunction with the revised codes, medical professionals will be able to detect and treat CKD in their patients earlier, the government will glean new information about the estimated 20 million people with CKD and related health problems such as diabetes, anemia and heart disease, and most importantly, the overall outcome of our patients with CKD will be optimal."

The change made by the NCHS resulted from a collaboration that included the NKF and the Renal Physicians Association. ICD-9-CM codes and classifies morbidity data from inpatient and outpatient records, physician offices, and most NCHS surveys. NCHS serves as the World Health Organization (WHO) Collaborating Center for the Family of International Classifications of North America. It is responsible for coordination of all official disease classification activities in the U.S. relating to the ICD and its use, interpretation, and periodic revision.

Other benefits of this revision of the ICD-9-CM codes include the ability to identify CKD patients who are also kidney transplant recipients, the ability to link specific treatments to the appropriate CKD stage, and unade toward achieving Healthy People 2010 goals.

For more information about the new codes, contact http://www.cdc.gov/nchs/datawh/ftpserv/?tpicd9/ftpicd9.htm#quidelines.

About the National Kidney Foundation

The National Kidney Foundation is a major voluntary health organization that seeks to prevent kidney and urinary tract diseases, improve the health and well-being of individuals and families affected by these diseases, and increase the availability of all organs for transplantation.

###

A GOOD AN INVESTMENT

The "Diabetes and Kidney Programs Line"

in the Department of Community Health budget:

- Preventing diabetes especially in children
- (kidney failure, blindness, amputation, heart attack, stroke) Preventing or delaying the complications of diabetes
 - Delaying kidney failure for those who already have chronic kidney disease 3.)

Diabetes and kidney disease prevention is possible, powerful and proven.

making wise food choices in order to reach and maintain a healthy body weight. Type 2 Diabetes can be prevented or delayed by exercising regularly and This is very important, especially in children.

Chronic kidney disease caused by diabetes can be prevented or delayed by keeping blood sugar levels and blood pressure as normal as possible. 🌯 EARLY Treatment No Treatment KIDNEY FAILURE Early treatment can extend kidney function in people with chronic kidney disease. 🖁

YEARS UNTIL KIDNEY FAILURE
Adopted from Brenner, et. al., 2001, courtesy of
National Kidney Disease Education Program



FACT

Clinical trials prove that diabetes and kidney disease prevention works.

-\\\

Prevention results show a fast turnaround in saving money.

PREVENTING CHRONIC KIDNEY DISEASE


National Kidney Of Michigan, Inc. **Foundation**™

IN THE HIGH-RISK AFRICAN AMERICAN POPULATION BY ADDRESSING CHRONIC KIDNEY DISEASE

12,191 beauty salon clients educated since 1999 "Healthy Hair Starts With a Healthy Body"



in the program. after participating 40% "prevention steps" Clients take



importance of regular check-ups Beauty salon stylists are trained to educate their kidney disease, nutrition, exercise and the clients about diabetes, high blood pressure,

"Dodge the Punch: Live Right" American barber shops to educate men. This is a new health campaign in African


"KICK" (Kids Interested in the Care of their Kidneys) 624,219 students since 1997 80% 20% 40% 60% "Kids & Kidneys" 87,170 kids since 2000


KIDNEY DISEASE PREVENTION, AND ORGAN DONATION BY TEACHING KIDS ABOUT IMPROVING THEIR HEALTH,

Students show a gain in knowledge after testing. What are the What is How much main causes hemodyalis? exercise is of kidney needed each disease? week?

"Healthy Kids & Kidneys"

- >Targets kids at high risk for diabetes or kidney disease >Builds upon school-based health clinics.
- >Provides individual and family counseling





Physical Activity Assessment

will focus on initiatives for children and adults, in communities and in health care settings, to prevent chronic kidney disease. Watch for GFR (% kidney function) results on your future laboratory blood tests! FUTURE: The newly released (March 8, 2005) Strategic Plan for Chronic Kidney Disease Prevention

Michigan Diabetes Outreach Network (MDON)

www.diabetesinmichigan.org 517-335-8445 Lansing, MI, Michigan Dept. of Community Health



National Kidney Foundation of Michigan (NKFM)

www.nkfm.org 800-482-1455 Ann Arbor MI 48108 1169 Oak Valley Drive




"Diabetes and Kidney Programs Line" implements effective programs with measurable outcomes.



PREVENTING COMPLICATIONS OF DIABETES

Michigan Diabetes Outreach Network

BY INCREASING TESTING FOR EARLY DETECTION



FUTURE: Preventing type 2 diabetes in children will be "a must" in future MDON programing in people's exercise habits increase by a 22% by a 14% increase in

43% decrease in amputations

21% decrease in death

complications preventing all

blindness

amputations preventing

heart disease

kidney disease preventing

and stroke

preventing

toot exams increase in

increase in

kidney testing increase in

lipid testing

preventing


Every 1% drop in A1c results in :

14% decrease in heart attacks 24% decrease in kidney failure

12% decrease in strokes

curriculums of the Michigan Model Programs For example: Incorporating programs like "Type 2 We're on to You" into health or science

Michigan Must Strengthen Efforts to Turn-Around **Epidemics of Diabetes and Kidney Disease**



Diabetes and chronic kidney disease are two of the most urgent health problems in the U.S. today.

Diabetes is reaching epidemic proportions. In the U.S. 16 million are afflicted. Children are at high risk because of obesity and inactivity.

More than 20 million Americans (1 in 9 adults) have chronic kidney disease, but most don't know it.

Diabetes caused 44% of kidney failure in 2003.

African Americans, Hispanics, Native Americans, and Asian Americans are all at high risk for diabetes, chronic kidney disease and kidney failure.


African Americans make up approximately 14% of Michigan's population but accounted for 46 % of the dialysis population in Michigan in 2002.

The increasing cost of health insurance threatens the business climate in Michigan.

Michigan Medicaid costs increase annually. (\$150 million in 2004)

Obesity and inactivity will cause 1 in 3 children born in 2000 to develop diabetes;

1 in 2 if African American or Hispanic.
-Center for Disease Control-



Fast-forward 20 years.......
Many of these "kids" will develop complications from diabetes in their 30's and 40's.